



E228

JACC March 27, 2012

Volume 59, Issue 13

**ACC-i2 with TCT**

CLINICAL OUTCOMES AFTER TRANSCATHETER AORTIC VALVE REPLACEMENT (TAVR) USING VALVE ACADEMIC RESEARCH CONSORTIUM (VARC) DEFINITIONS: A WEIGHTED META-ANALYSIS OF 3,519 PATIENTS FROM 17 STUDIES

i2 Poster Contributions

McCormick Place South, Hall A

Saturday, March 24, 2012, 9:30 a.m.-Noon

Session Title: Outcomes in Patients Treated with TAVR

Abstract Category: 20. PCI - Valvular Heart Disease

Presentation Number: 2533-233

Authors: *Philippe Genereux, Stuart Head, Nicolas Van Mieghem, Susheel Kodali, Ajay Kirtane, Ke Xu, Jeffrey Moses, Craig Smith, Patrick Serruys, A. Kappetein, Martin Leon, Columbia University Medical Center and The Cardiovascular Research Foundation, New York, NY, USA, Hôpital du Sacré-Coeur de Montréal, Université de Montréal, Montréal, Canada*

Backgrounds: Recently, the published Valve Academic Research Consortium (VARC) definitions have helped to add uniformity for reporting outcomes after Transcatheter Aortic Valve Replacement (TAVR). We sought to perform a weighted meta-analysis to determine rates of major outcomes after TAVR using VARC definitions and to evaluate their current use in the literature.

Methods: A comprehensive search of multiple electronic databases from January 1st 2011 through October 12th 2011 was conducted using predefined criteria. We included studies reporting at least one outcome using VARC definitions.

Results: A total of 17 studies including 3,519 patients met inclusion criteria and were included in the analysis. The pooled estimate rate of outcomes were determined according to VARC's definitions: device success: 92.1%, 95%CI [88.7,95.5]; all cause 30-day mortality: 7.8%, 95%CI [5.5,11.1]; myocardial infarction: 1.1%, 95% CI [0.2,2.0]; acute kidney injury stage II-III: 7.5%, 95%CI [5.1,11.4]; life threatening bleeding: 15.6%, 95% CI [11.7,20.7]; major vascular complication: 11.9%, 95% CI [8.6,16.4]; major stroke 3.2%, 95%CI [2.1,4.8]; and new permanent pace maker (PPM) implantation: 13.9%, 95% CI [10.6,18.9]. Medtronic Corevalve™ prosthesis use was associated with a significant higher rate of PPM implantation compared to the Edwards's prosthesis (28.9%, 95% CI [23.0,36.0] vs. 4.9%, 95% CI [3.9,6.2], p value < 0.0001). The 30-day safety composite endpoint rate was 32.7%, 95%CI [27.5,38.8] and the 1-year total mortality was 22.1%, 95% CI [17.9,26.9].

Conclusion: VARC definitions have already been used by the TAVR clinical research community, establishing a new standard for reporting clinical outcomes. Future revisions of the VARC definitions are needed based upon evolving TAVR clinical experiences.